


Vol. XXVI - No. 1 - January/March 2011 - Trimestrale - ISSN 0393 - 5264

In caso di mancato recapito, rinviare a Roma Romanina Stampe, per restituzione al mittente previo pagamento resi



Istituto di Ricovero e Cura
a Carattere Scientifico
Fondazione
"Istituto Neurologico
C. Mondino" - Pavia

FUNCTIONAL NEUROLOGY



EDITOR-IN-CHIEF
STEFANO BASTIANELLO

Quarterly published by



CIC Edizioni Internazionali

www.functionalneurology.it

functional neurology

and new trends in interventional neurosciences

Editor-in-Chief

STEFANO BASTIANELLO (*Pavia, Rome - Italy*)

Founding Editor

GIUSEPPE NAPPI (*Pavia, Rome - Italy*)

Senior Editors

PIERFRANCO SPANO (*Brescia - Italy*)

OTTAR SJAASTAD (*Trondheim - Norway*)

Associate Editor

EGIDIO D'ANGELO (*Pavia - Italy*)

Editorial Assistants

SILVIA MOLINARI (*Pavia - Italy*)

E-mail: silvia.molinari@mondino.it

GUENDALINA FORNARI LUSWERGH (*Rome - Italy*)

Language Editor

CATHERINE WRENN (*Pavia - Italy*)

Editorial Office

IRCCS Fondazione "Istituto Neurologico C. Mondino"

Via Mondino, 2 - 27100 Pavia, Italy

Tel. +39 0382 380299 - Fax +39 0382 380448

E-mail: funcneur@mondino.it

Publisher

CIC Edizioni Internazionali srl

Corso Trieste 42

00198 Rome, Italy

Tel. +39 06 8412673

Fax +39 06 8412688

E-mail: info@gruppocic.it

www.gruppocic.it

Managing Editor

(Direttore Responsabile)

ANDREA SALVATI

Rome, Italy

E-mail: a.salvati@gruppocic.it

Managing Office

ANTONELLA ONORI

Rome, Italy

E-mail: onori@gruppocic.it

Advertising Office

PATRIZIA ARCANGIOLI

Rome, Italy

E-mail: arcangioli@gruppocic.it

Consulting Editors**Neuroanatomy**

Michele Papa (Italy), Rafi Malach (Israel)

Neurochemistry and neurobiology

Fabio Blandini (Pavia - Italy), J. Timothy Greenamyre (Pittsburgh - USA)

Neurodegenerative diseases

Emilia Martignoni (Varese - Italy), Amos D. Korczyn (Ramat Aviv - Israel)

Neuroendocrinology

Rossella E. Nappi (Pavia - Italy), Ashley B. Grossman (London - UK)

Neuroepidemiology

Antonietta Citterio (Pavia - Italy), W. Allen Hauser (New York - USA)

Neuroesthetics

Klaus Podoll (Aquisgrana - Germany), Ubaldo Nicola (Pavia - Italy)

Neuroethics

Francesco Maria Avato (Ferrara - Italy), Livio P. Tronconi (Pavia - Italy)

Neurogenetics

Filippo M. Santorelli (Rome - Italy), Giovanni Stevanin (Paris - France)

Neurogenic pain and headache

Cristina Tassorelli (Pavia - Italy), Rigmor H. Jensen (Copenhagen - Denmark)

Neurofunctional imaging

Richard Frackowiack (London - UK), Gian Luigi Lenzi (Rome - Italy)

Neuromolecular and cellular imaging

Jeff Bulte (Baltimore - USA), Andreas H. Jacobs (Cologne - Germany)

Neuro-oncology

Felice Giangaspero (Rome - Italy), Robert L. Martuza (Boston - USA)

Neuropharmacology

Maurizio Memo (Brescia - Italy), Emilio Perucca (Pavia - Italy)

Neurophilosophy

Michele Di Francesco (Milan - Italy), Patricia Smith Churchland (San Diego - USA)

Neurophysiology

Arrigo Moglia (Pavia - Italy), Trond Sand (Trondheim - Norway)

Neuropsychology

Carlo Caltagirone (Rome - Italy), François Boller (Bethesda - USA)

Neuroradiology

Rudiger von Kummer (Dresden - Germany), Cesare Colosimo (Rome - Italy)

Neurorehabilitation

Giorgio Sandrini (Pavia - Italy), Leopold Saltuari (Innsbruck - Austria)

Neuroscience history

Luciana R. Angeletti (Rome - Italy), Paolo Mazzarello (Pavia - Italy)

Neurosurgery

Roberto Delfini (Rome - Italy), Francesco Tomasello (Messina - Italy)

Neurotoxicology

Laura Berliocchi (London - UK), Oliver J. Dolly (Dublin - Ireland)

Neurovascular disorders

Antonio Carolei (Aquila - Italy), J. Donald Easton (Providence RI - USA)

Neurovenous disorders

Paolo Zamboni (Italy), Robert Zivadinov (Buffalo - USA)

Contents

COMA AND CONSCIOUSNESS

Guest editor: Michele Papa (Naples, Italy)

EDITORIAL

WHAT A HUMAN BEING IS

A. Papa-Grimaldi

5

*OSCILLATORY BRAIN ACTIVITY IN
VEGETATIVE AND MINIMALLY CONSCIOUS
STATE DURING A SENTENCE
COMPREHENSION TASK*

M. Schabus, C. Pelikan, N. Chwala-Schlegel,
K. Weilhart, D. Roehm, J. Donis, G. Michitsch,
G. Pichler, W. Klimesch

31

ORIGINAL ARTICLES

*THE ROLE OF EARLY ELECTROCLINICAL
ASSESSMENT IN IMPROVING
THE EVALUATION OF PATIENTS WITH
DISORDERS OF CONSCIOUSNESS*

A. Amantini, R. Carrai, S. Fossi, F. Pinto,
A. Grippo

7

*RESTING STATE ACTIVITY IN PATIENTS
WITH DISORDERS OF CONSCIOUSNESS*

A. Soddu, A. Vanhaudenhuyse, A. Demertzi,
M.-A. Bruno, L. Tshibanda, H. Di, M. Boly,
M. Papa, S. Laureys, Q. Noirhomme

37

*VEGETATIVE STATE, MINIMALLY
CONSCIOUS STATE, AKINETIC MUTISM
AND PARKINSONISM AS A CONTINUUM
OF RECOVERY FROM DISORDERS OF
CONSCIOUSNESS: AN EXPLORATORY
AND PRELIMINARY STUDY*

R. Formisano, M. D'Ippolito, M. Risetti,
A. Riccio, C. Falletta Caravasso, S. Catani,
F. Rizza, A. Forcina, M.G. Buzzi

15

*POST-ANOXIC VEGETATIVE STATE:
IMAGING AND PROGNOSTIC
PERSPECTIVES*

M. Stanziano, C. Foglia, A. Soddu,
F. Gargano, M. Papa

45

*DISORDERS OF CONSCIOUSNESS AND
COMMUNICATION. ETHICAL MOTIVATIONS
AND COMMUNICATION-ENABLING
ATTRIBUTES OF CONSCIOUSNESS*

G. Tamburrini, D. Mattia

51

*AUTOMATED EEG ENTROPY
MEASUREMENTS IN COMA, VEGETATIVE
STATE/UNRESPONSIVE WAKEFULNESS
SYNDROME AND MINIMALLY
CONSCIOUS STATE*

O. Gosseries, C. Schnakers, D. Ledoux,
A. Vanhaudenhuyse, M.-A. Bruno,
A. Demertzi, Q. Noirhomme, R. Lehembre,
P. Damas, S. Goldman, E. Peeters,
G. Moonen, S. Laureys

25,

*COGNITIVE REHABILITATION
IN NON-COMMUNICATIVE
BRAIN-DAMAGED PATIENTS*

L. Trojano, P. Moretta, A. Cozzolino,
A. Saltalamacchia, A. Estraneo

55

FORTHCOMING EVENTS

63

Post-anoxic vegetative state: imaging and prognostic perspectives

Mario Stanziano^a
Carolina Foglia, MD^a
Andrea Soddu, PhD^{a,b}
Francesca Gargano, MD^a
Michele Papa, MD^a

^a Laboratorio di Morfologia delle Reti Neuronalì,
Department of Public, Clinical and Preventive
Medicine, Second University of Naples, Italy

^b Coma Science Group, Cyclotron Research Centre,
University of Liège, Belgium

Corresponding author: Mario Stanziano
Laboratorio di Morfologia delle Reti Neuronalì
Dipartimento Medicina Pubblica Clinica e Preventiva
Second University of Naples
Via L. Armanni 5, 80138 Naples, Italy
E-mail: mariostanziano@libero.it

Summary

Prognostic determination of patients in coma after resuscitation from cardiac arrest is a common and difficult requirement with significant ethical, social and legal implications.

We set out to seek markers that can be used for the early detection of patients with a poor prognosis, so as to reduce uncertainty over treatment and non-treatment decisions, and to improve relationships with families.

We reviewed the medical literature from 1991 to 2010, using key words such as post-anoxic coma, post-anoxic vegetative state, vegetative state prognosis, recovery after cardiac arrest.

Neurological examination, electrophysiology, imaging, and biochemical markers are all useful tools for estimating patients' chances of recovery from cardiac arrest.

It seems unlikely that any single test will prove to have 100% predictive value for outcome; but the combination of various prognostic markers, as shown in some articles, could increase the reliability of outcome prediction. However, further research is needed.

KEY WORDS: anoxic brain injury, prognostic factors, vegetative state

Introduction

Anoxia is the third most frequent cause of coma, after trauma and vascular lesions. The most common causes of post-anoxic coma in adults are: cardiopulmonary arrest, stroke, respiratory arrest and carbon monoxide poisoning (1). In children, the causes include asphyxia, congenital malformations, and perinatal injuries (2). The incidence of hypoxic-ischemic brain injury is not well known, but it is certain that cardiac arrest, the most com-

mon cause of post-anoxic coma, affects about 450000 Americans per year (3). Almost eighty per cent of patients who initially survive a cardiac arrest remain comatose for varying lengths of time, and approximately forty per cent of these progress to vegetative state (4). Early identification of patients with a poor prognosis is desirable in order to reduce uncertainty over treatment and non-treatment decisions, and to improve relationships with families. Various clinical parameters, neurological examination models, biochemical tests, neuroimaging and electrophysiological techniques have been proposed for the prognostic evaluation of brain function in comatose cardiac arrest survivors.

Clinical-prognostic factors

Some factors, such as age, duration of arrest and duration of coma, have been investigated as predictors of functional outcome in post-anoxic patients. To consider age a negative prognostic factor is still controversial. Schultz et al. showed a significant difference in survival between patients under 60 and those over 80 years (15% vs 4%, respectively) (5) and Roest et al. observed that prognosis is twice as good in younger patients not only at 30 days but also at 180 days after coma onset (6). Conversely, Berger and Kelley in a prospective analysis of 255 in-hospital cardiopulmonary arrests in non-critical patients demonstrated that age was not an independent predictor of survival (7). In addition, Rogove et al., in a prospective study of 774 patients, found that old age did not negate good cerebral outcome after cardiorespiratory arrest (8). In reality, to make an overall assessment, we should also take into account patients' comorbidities and complications of cardiac arrest. Since the duration of cardiac arrest correlates with the extent of brain damage, one can expect to observe a direct proportional relationship between the duration of the arrest and the outcome (8,9). Several studies have indeed shown a close relationship: the survival rate has been found to be 48% for a period of less than 10 minutes and 2% for longer than 10 minutes (5). Saklayen et al. confirmed that a shorter duration of anoxia correlates with a better prognosis (10), but in practice it is difficult to determine the duration of arrest when evaluating a patient. As noted before, up to 80% of patients are comatose after an arrest. Of those patients destined to awaken, over 90% will do so within the first 72 hours (10). Thereafter, the probability of awakening decreases progressively. In fact, in post-anoxic coma, the probability of recovery after three months of vegetative state becomes statistically close to zero, and at this point the condition is defined permanent vegetative state (11). The most widely studied and reliable predictors of neurological outcome continue to be provided

by the bedside neurological examination (12,13). An absent pupillary reaction to light after 72 hours carries a prognosis of coma until death with 100% specificity (14,15). Another parameter widely discussed is the motor response to painful stimuli. Absence of this response after three days of coma is related to a poor outcome with a specificity of 100%, although not all patients with poor outcomes were identified only by this parameter (16). Moreover, in the Brain Resuscitation Clinical Trials I, lack of the motor response after 72 hours was the only independent predictor of poor outcome (17). The corneal reflex is tested by touching the cornea with a gauze or cotton swab and looking for contraction of the orbicularis oculi on either side. In two prospective studies, absence of the corneal reflex at 72 hours was associated with no false positives for a poor outcome (18,19). The post-resuscitation clinical course has also been found to influence the outcome of these patients. The occurrence of status epilepticus in cardiac arrest survivors has been strongly associated with mortality (20). Many studies have also considered myoclonic status epilepticus (bilaterally synchronous twitches of limb, trunk, or facial muscles) as a predictor of poor outcome in patients with severe injury (19,21). In a prospective study involving 407 patients, myoclonic status epilepticus at 24 hours after cardiac arrest was associated with no false positives (19). The Glasgow Coma Scale (GCS) score has been extensively investigated as a predictor of individual outcome following cardiac arrest with cardiopulmonary resuscitation. In particular, the Cerebral Resuscitation Study Group of the Belgian Society for Intensive Care found GCS scores of <5 and >9 to predict poor and good outcome respectively. For patients with GCS scores of 5 to 9, a repeated score >8 during the first six days was shown to be a cut-off for good prognosis (22). Others claimed that a GCS value of <4 after three days of coma was the best indicator of poor outcome with a specificity of 100% (17). Zandbergen et al., in a systematic review, showed that a GCS score of 5 or less in the first 24 hours was not helpful in predicting outcome (14). Results of a recent study, conducted in patients treated with hypothermia, showed that at day 3 after cardiac arrest a GCS score >6 had a specificity of 87% and a sensitivity of 73% for predicting a good outcome (23). In view of these (to date) conflicting data, the GCS cannot be considered a sufficiently sensitive measure compared with other tests of central nervous system function (24). Fugate et al. conducted a study to determine whether the FOUR score is an accurate predictor of outcome in patients after cardiac arrest. They showed that most patients with a FOUR score >8 at days 3-5 after cardiac arrest survived to hospital discharge, while no patient with a FOUR score <4 survived (25). In the future, this simple score could be used as a prognostic tool, but further studies are needed to confirm its validity.

Neurophysiological prognostic factors

The electroencephalogram (EEG) during cardiac arrest is isoelectric. Half of all patients, immediately after resuscitation, do not show cortical activity on EEG (15). This finding does not reliably predict poor outcome (15) and it is usual for clinicians to wait at least 24 hours be-

fore ordering an EEG in order to avoid falsely disappointing results. Several unequivocal patterns have prognostic value after this time. In particular, a generalized suppression $\leq 20 \mu\text{V}$, burst suppression with generalized epileptiform activity, and generalized periodic complexes on a flat background are related to poor outcome (6). In combining data from different trials, Zandbergen et al. found that an EEG with an isoelectric or burst suppression pattern during the first week after resuscitation had a 100% positive predictive value for coma until death or persistent vegetative state (14). Recently, some authors have remarked on the prognostic significance of reactivity in alpha-theta coma (ATC), either at the first observation or during the evolution of the electroencephalographic pattern, suggesting that there exist two forms of ATC having different prognostic value. They identified a complete ATC with a-reactive, monotonous, continuous and frontally distributed alpha activity, typically associated with abnormal or absent cortical somatosensory evoked potentials (SEPs). This variant is associated with little improvement in the first 48 hours after cardiac arrest and poor outcome (26). The other form is incomplete ATC, characterized by less monotonous, posteriorly accentuated and partially reactive EEG activity typically associated with normal SEPs. It is associated with a better prognosis and with the possibility of recovery of consciousness (26,27). Sensory evoked potentials, especially SEPs, have been investigated for their potential to predict long term outcome from coma (over 20 years) (28). SEPs are non-invasive, reproducible, and simple to perform and to interpret. In addition, compared with the EEG, they are less susceptible to electrical interference and less affected by metabolic disturbances or medications, although it is recommended to wait 24 hours before measuring them (29). They test the integrity of the neuronal pathways from the peripheral nerve, spinal cord, brainstem and cerebral cortex. The N20 component, which is the first cortical response of the SEP, is the evoked potential waveform best studied as a prognostic indicator. Absence of N20, after stimulation of the median nerve, has been shown to be a reliable predictor of poor outcome in several studies of patients in coma after cardiac arrest (19,29-34). Recently, a systematic review of 18 studies analyzed the predictive ability of SEPs acquired early after onset of coma in 1136 adult patients with hypoxic-ischemic encephalopathy. It was found that all 336 patients with bilaterally absent cortical N20 SEP peaks did not awake from coma (35). Two other clinical trials also showed that absence of cortical SEP peaks was associated with poor prognosis, with a specificity of 100% (36,37). However, the presence of cortical N20 SEP responses is not a guarantee for awakening from coma (36). To improve the accuracy of SEPs in predicting good outcomes in comatose cardiac arrest survivors, Madl and coworkers evaluated long-latency N70 peaks in 66 patients. In this series, N70 latency was less than 118 milliseconds in all patients who had a good recovery and was absent or greater than 118 milliseconds in patients who had a poor outcome (38). In recent years, it has been suggested that routine clinical tests should also include the assessment of higher-order cortical activity via endogenous evoked potentials (EPs). The presence of a P300 component in comatose patients proved to be effective in predicting awakening in small patient series,

while its absence does not necessarily predict a good prognosis (39,40). However, since the P300, which is agreed to be associated with attention or expectancy, is dependent on the level of vigilance, it is preferable to record mismatch negativity (MMN), which does not require awareness. Furthermore, several studies have shown that the presence of MMN in a comatose patient is associated with a high percentage of evolution toward awakening (41-43). In summary, it can be concluded that absent exogenous EPs are well established prognosticators of poor outcome, whereas the presence of endogenous components, notably the MMN and P300, appears to predict a favorable outcome (44).

Biochemical prognostic markers

In recent years it has been shown that increased serum levels of neuron-specific enolase (NSE) and of the astroglial protein S100 are associated with hypoxic-ischemic brain injury and unfavorable neurological outcome (45). S100 is a calcium-binding protein present in high concentrations in glial and Schwann cells. Serum levels of S100 increase transiently, reaching a peak within 24 hours of the anoxic episode, and continuing to show high, but declining, values on the following two days. This initial rise may reflect reversible brain edema combined with a disturbance of astroglial cell membrane integrity and blood-brain barrier function. The low weight and high solubility of the molecule probably facilitate this process. In cases with a short period of anoxia, the pathological S100 levels rapidly return to normal within 24 hours, which is compatible with the fast turnover of S100 in serum. Values remaining high or increasing on days 2 and 3 may indicate permanent brain damage, which is of importance for the prognosis (46,47). Moreover, levels elevated above 0.217 $\mu\text{g/l}$ on day 2 after return of spontaneous circulation indicate severe neurological injury and predict poor outcome with a 100% positive predictive value (47). Another study demonstrated that at 48 hours after cardiac arrest, an S100 serum level $>1.10 \mu\text{g/l}$ has a specificity of 100% for the diagnosis of brain damage (48). However, both studies, despite the different cut-off, claim that S100 represents an early marker of short-term outcome after cardiac arrest. NSE is the neuronal form of the intracytoplasmic glycolytic enzyme enolase. As opposed to the transient increase in S100, levels of NSE have been found to increase continuously after cardiac arrest and to peak on the third day. This is in line with the fact that NSE has shown promising results, in terms of prognosis, at later stages after cardiac arrest (48-51). Although cerebrospinal fluid (CSF) levels of NSE are more sensitive for neuronal damage, a serum level above 23.2 $\mu\text{g/l}$ has a 100% positive predictive value for coma, vegetative state, or death (41). In one prospective multicenter study involving 231 patients, an NSE level of more than 33 $\mu\text{g/l}$, sampled between 1 and 3 days after cardiac arrest, was strongly predictive of a poor outcome with no false positives (19,51). Instead, elevated levels of S100, as indeed the BB fraction of creatine kinase in CSF or serum and the neurofilament protein, did not show high specificity for a poor outcome (51). It is worth noting that the widespread use of biochemical markers is limited due to widely varying cut-off levels for

identifying specific outcomes, inconsistency in sampling times, and lack of uniformity in patient populations.

Neuroimaging prognostic factors

Computed tomographic (CT) images are usually normal immediately after a cardiac arrest, but by day 3 they often show brain swelling and reversal of the gray/white matter densities (with the use of quantitative measures) in patients with a poor outcome (35). Further study is needed to assess the clinical use of these findings in establishing prognosis. Magnetic resonance imaging (MRI) has also been proposed as a means of assessing prognosis after cardiac arrest, but limited data call its use into question. Among four case series, each with no more than 12 patients assessed with MRI at variable time points (52,53), two studies showed that diffusely abnormal findings on diffusion weighted imaging (DWI) and fluid-attenuated inversion recovery (FLAIR) correlated with a poor outcome (54,55). DWI detects cytotoxic edema by measuring the random motion of water protons, a process that is reduced by failure of the energy-requiring active water transport mechanism. DWI studies in animal models have demonstrated a decline in brain apparent diffusion coefficient (ADC) values during cardiac arrest, which then reverse after successful resuscitation (56,57). Despite successful reperfusion, however, secondary energy failure and ADC decrease appear after several hours (58). It has been shown in preliminary studies that quantitative MRI brain changes are correlated with functional outcome in comatose cardiac arrest survivors (59,60). The percentage of brain tissue below a threshold of $650 \times 10^{-6} \text{ mm}^2/\text{s}$ to $700 \times 10^{-6} \text{ mm}^2/\text{s}$ was found to be correlated with functional outcome at three months after the arrest (59). Based on whole-brain quantitative DWI analyses, the ideal time window for prognostication appears to be between 49 and 108 hours after the arrest, when the ADC reductions are most apparent. None of the patients with $>10\%$ of brain tissue with an ADC value $<650 \times 10^{-6} \text{ mm}^2/\text{s}$ to $700 \times 10^{-6} \text{ mm}^2/\text{s}$ during this time window regained consciousness. In a prospective study, Mlynash et al. analyzed in detail which brain structures at which time point were the most severely affected by ADC reductions by comparing good- and poor-outcome patients with normal controls and with each other (61). In agreement with the results of another retrospective study (60), they found that ADC changes, in post-cardiac arrest patients, are both time- and region-dependent during the first week. In addition, both the qualitative and the quantitative MRI changes in poor-outcome patients were most severe in the cortical regions and most apparent between three and five days after the arrest. Indeed, in these conditions, although ADC changes occur globally, they most profoundly affect the cortical gray matter structures. By contrast, patients who are able to wake up from their coma exhibit increased diffusion involving the temporal and occipital lobes, corona radiata, and hippocampus; in addition, qualitative changes in the deep gray nuclei alone are common in these patients. MR spectroscopy (e.g., for pH and N-acetyl aspartate, a neuronal marker) has been reported to correlate with a poor outcome in small studies (60,62), but more data are needed. Functional brain imaging in comatose sur-

vivors of cardiac arrest could provide important diagnostic and prognostic information beyond that available from bedside examination alone (63,64). Di et al. reviewed 15 positron emission tomography (PET) and functional magnetic resonance imaging (fMRI) studies involving 48 published cases which were classified as 'absent cortical activation', 'typical activation' (involving low-level primary sensory cortices), and 'atypical activation' (corresponding to higher-level associative cortices). The results show that atypical patterns of activity appear to predict recovery from the vegetative state with 93% specificity and 69% sensitivity. That is to say, nine of 11 (3 post-anoxic) patients exhibiting atypical patterns of activity recovered consciousness, whereas 21 of 25 (18 post-anoxic) patients with typical primary cortical activity patterns, and four out of four (3 post-anoxic) patients with absent activity failed to recover (65). In another recent study of 41 patients with disorders of consciousness, Coleman et al. found direct evidence of prognostically important information within neuroimaging data that was at odds with the behavioral assessment performed at the time of scanning. Thus, contrary to the clinical impression of a specialist team using behavioral assessment tools, two patients (one with brainstem stroke) who had been referred to the study with a diagnosis of vegetative state did in fact demonstrate clear signs of speech comprehension when assessed using fMRI. More importantly, however, across the whole group of patients, the fMRI data were found to have no association with the behavioral presentation at the time of the investigation, but correlated significantly with subsequent behavioral recovery, six months after the scan. In this case, the fMRI data predicted subsequent recovery in a way that a specialist behavioral assessment could not (66). Another interesting study is the one proposed by Gofton et al. in which, using fMRI, the authors measured cerebral activation in response to somatosensory stimulation of the palm of the hand in 19 comatose survivors of cardiac arrest and in 10 healthy control subjects and compared it to SEP testing of the median nerve. They demonstrated that patients with a favorable outcome and with SEP responses had a greater blood oxygenation-level dependent (BOLD) signal in the primary somatosensory cortex (S1) contralateral to the stimulated hand, when compared with patients with an unfavorable outcome and without SEP responses. Furthermore, there also emerged positive correlations between BOLD in S1 and both levels of consciousness and clinical measures of outcome at three months (67). However, further studies should be done before fMRI can be deemed a useful tool for predicting outcome of patients in post-anoxic coma. Most of the studies on prognostic markers in cardiac arrest have used single variables. However, several studies (68,69) have looked at combining the values of the above tests to increase the reliability of outcome predictions. One example is the study by Bassetti and colleagues that assessed the value of combining clinical examination, EEG, SEPs, and two serum biochemical markers (ionized calcium and NSE). Individually, clinical examination correctly predicted outcome in 58%, SEPs in 59%, EEG in 41%, while combining the three factors (clinical examination, SEPs, and EEG) increased the rate of correct predictions to 82%. No false pessimistic prediction was observed using the combination (69).

Concluding remarks

From this review of the literature, there emerges a series of key points in relation to the prognosis of patients in post-anoxic coma. First of all, **age** is not a significant factor in outcome, especially when considered alone; conversely, **cardiac arrest lasting** more than 10 minutes and **coma lasting** more than 24 hours are good individual predictors of poor outcome. Absence of a **pupillary reaction** to light, absence of the **corneal reflex** and absence of a **motor response** to painful stimuli at 72 hours are each suggestive of a poor outcome. Also indicative of a poor outcome is an **EEG** with an isoelectric or burst suppression pattern during the first week after resuscitation and/or a complete ATC.

Somatosensory evoked potentials have shown some usefulness and are unaffected by confounding factors. Abnormalities are 100% specific for a poor outcome, although only after at least 24 hours of coma. The absence of abnormalities, on the other hand, does not necessarily predict a good outcome.

The predictive value of **neurobiochemical markers** is variable; these markers are unhelpful before 48 hours have elapsed.

Neuroimaging after return of spontaneous circulation has not been studied extensively but recent data indicate that a diffuse cortical hypersignal on DWI may predict a poor outcome. It is possible that fMRI, in the near future, will become very helpful in predicting outcome in comatose patients.

Finally, **combining** various prognostic markers could increase outcome prediction reliability.

References

1. Tresch DD, Sims FH, Duthie EH, Goldstein MD, Lane PS. Clinical characteristics of patients in the persistent vegetative state. *Arch Neurol* 1991;151:930-932
2. Ashwal S, Bale JF Jr, Coulter DL et al. The persistent vegetative state in children: report of the Child Neurology Society Ethics Committee. *Ann Neurol* 1992;32:570-576
3. Callans DJ. Out-of-hospital cardiac arrest—the solution is shocking. *N Engl J Med* 2004;351:632-634
4. Madl C, Holzer M. Brain function after resuscitation from cardiac arrest. *Curr Opin Crit Care* 2004;10:213-217
5. Schultz SC, Cullinane DC, Pasquale MD, Magnant C, Evans SR. Predicting in-hospital mortality during cardiopulmonary resuscitation. *Resuscitation* 1996;33:13-17
6. Roest A, van Bets B, Jorens PG, Baar I, Weyler J, Mercelis R. The prognostic value of the EEG in postanoxic coma. *Neurocrit Care* 2009;10:318-325
7. Berger R, Kelley M. Survival after in-hospital cardiopulmonary arrest of noncritically ill patients. A prospective study. *Chest* 1994;106:872-879
8. Rogove HJ, Safar P, Sutton-Tyrrell K, Abramson NS. Old age does not negate good cerebral outcome after cardiopulmonary resuscitation: analyses from the brain resuscitation clinical trials. The Brain Resuscitation Clinical Trial I and II Study Groups. *Crit Care Med* 1995;23:18-25
9. Grubb NR. Managing out-of-hospital cardiac arrest survivors: 1. Neurological perspective. *Heart* 2001;85:6-8
10. Saklayen M, Liss H, Markert R. In-hospital cardiopulmonary resuscitation. Survival in 1 hospital and literature review. *Medicine (Baltimore)* 1995;74:163-175
11. Medical aspects of the persistent vegetative state (1). The

- Multi-Society Task Force on PVS. *N Engl J Med* 1994;330:1499-1508
12. Pütgen HA, Geocadin R. Predicting neurological outcome following cardiac arrest. *J Neurol Sci* 2007;261:108-117
 13. Young GB. Clinical practice. Neurologic prognosis after cardiac arrest. *N Engl J Med* 2009;361:605-611
 14. Zandbergen EG, de Haan RJ, Stoutenbeek CP, Koelman JH, Hijdra A. Systematic review of early prediction of poor outcome in anoxic-ischaemic coma. *Lancet* 1998;352:1808-1812
 15. Jørgensen EO, Holm S. Prediction of neurological outcome after cardiopulmonary resuscitation. *Resuscitation* 1999;41:145-152
 16. Codazzi D, Pifferi S, Savioli M, Langer M. Neurologic prognosis after cardiocirculatory arrest outside the hospital. [Article in Italian] *Minerva Anestesiol* 1997;63:353-364
 17. Edgren E, Hedstrand U, Kelsey S, Sutton-Tyrrell K, Safar P. Assessment of neurological prognosis in comatose survivors of cardiac arrest. BRCT I Study Group. *Lancet* 1994;343:1055-1059
 18. Berek K, Lechleitner P, Luef G et al. Early determination of neurological outcome after prehospital cardiac arrest. *Stroke* 1995;26:543-549
 19. Zandbergen EG, Hijdra A, Koelman JH et al.; PROPAC Study Group. Prediction of poor outcome within the first 3 days of postanoxic coma. *Neurology* 2006;66:62-68
 20. Rossetti AO, Logroscino G, Liaudet L et al. Status epilepticus: an independent outcome predictor after cerebral anoxia. *Neurology* 2007;69:255-260
 21. Wijdicks EF, Parisi JE, Sharbrough FW. Prognostic value of myoclonus status in comatose survivors of cardiac arrest. *Ann Neurol* 1994;35:239-243
 22. Mullie A, Verstringe P, Buylaert W et al. Predictive value of Glasgow Coma Score for awakening after out-of-hospital cardiac arrest. Cerebral Resuscitation Study Group of the Belgian Society for Intensive Care. *Lancet* 1988;1:137-140
 23. Scheffold JC, Storm C, Krüger A, Ploner CJ, Hasper D. The Glasgow Coma Score is a predictor of good outcome in cardiac arrest patients treated with therapeutic hypothermia. *Resuscitation* 2009;80:658-661
 24. Kaye P. Early prediction of individual outcome following cardiopulmonary resuscitation: systematic review. *Emerg Med J* 2005;22:700-705
 25. Fugate JE, Rabinstein AA, Claassen DO, White RD, Wijdicks EF. The FOUR score predicts outcome in patients after cardiac arrest. *Neurocrit Care* 2010;13:205-210
 26. Berkhoff M, Donati F, Bassetti C. Postanoxic alpha (theta) coma: a reappraisal of its prognostic significance. *Clin Neurophysiol* 2000;111:297-304
 27. Fossi S, Amantini A, Grippo A, Cossu C, Boni N, Pinto F. Anoxic-ischemic alpha coma: prognostic significance of the incomplete variant. *Neurol Sci* 2004;24:397-400
 28. Ahmed I. Use of somatosensory evoked responses in the prediction of outcome from coma. *Clin Electroencephalogr* 1988;19:78-86
 29. Zandbergen EG, de Haan RJ, Koelman JH, Hijdra A. Prediction of poor outcome in anoxic-ischemic coma. *J Clin Neurophysiol* 2000;17:498-501
 30. Kirsch M, Boveroux P, Massion P et al. Predicting prognosis in post-anoxic coma [Article in French]. *Rev Med Liege* 2008;63:263-268
 31. Koenig MA, Kaplan PW, Thakor NV. Clinical neurophysiologic monitoring and brain injury from cardiac arrest. *Neurol Clin* 2006;24:89-106
 32. Geocadin RG, Eleff SM. Cardiac arrest resuscitation: neurologic prognostication and brain death. *Curr Opin Crit Care* 2008;14:261-268
 33. Young GB, Doig G, Ragazzoni A. Anoxic-ischemic encephalopathy: clinical and electrophysiological associations with outcome. *Neurocrit Care* 2005;2:159-164
 34. Rothstein TL. The utility of median somatosensory evoked potentials in anoxic-ischemic coma. *Rev Neurosci* 2009;20:221-233
 35. Robinson LR, Micklesen PJ, Tirschwell DL, Lew HL. Predictive value of somatosensory evoked potentials for awakening from coma. *Crit Care Med* 2003;31:960-967
 36. Logi F, Fischer C, Murri L, Mauguière F. The prognostic value of evoked responses from primary somatosensory and auditory cortex in comatose patients. *Clin Neurophysiol* 2003;114:1615-1627
 37. Zingler VC, Krumm B, Bertsch T, Fassbender K, Pohlmann-Eden B. Early prediction of neurological outcome after cardiopulmonary resuscitation: a multimodal approach combining neurobiochemical and electrophysiological investigations may provide high prognostic certainty in patients after cardiac arrest. *Eur Neurol* 2003;49:79-84
 38. Madl C, Kramer L, Domanovits H et al. Improved outcome prediction in unconscious cardiac arrest survivors with sensory evoked potentials compared with clinical assessment. *Crit Care Med* 2000;28:721-726
 39. Yingling CD, Hosobuchi Y, Harrington M. P300 as a predictor of recovery from coma. *Lancet* 1990;336:873
 40. De Giorgio CM, Rabinowicz AL, Gott PS. Predictive value of P300 event-related potentials compared with EEG and somatosensory evoked potentials in non-traumatic coma. *Acta Neurol Scand* 1993;87:423-442
 41. Fischer C, Luaute J, Némoz C, Morlet D, Kirkorian G, Mauguière F. Improved prediction of awakening or nonawakening from severe anoxic coma using tree-based classification analysis. *Crit Care Med* 2006;34:1520-1534
 42. Fischer C, Luaute J, Adeleine P, Morlet D. Predictive value of sensory and cognitive evoked potentials for awakening from coma. *Neurology* 2004;63:669-673
 43. Fischer C, Morlet D, Bouchet P, Luaute J, Jourdan C, Salord F. Mismatch negativity and late auditory evoked potentials in comatose patients. *Clin Neurophysiol* 1999;110:1601-1610
 44. Daltrozzo J, Wioland N, Mutschler V, Kotchoubey B. Predicting coma and other low responsive patients outcome using event-related brain potentials: a meta-analysis. *Clin Neurophysiol* 2007;118:606-614
 45. Snyder-Ramos SA, Böttiger BW. Molecular markers of brain damage – clinical and ethical implications with particular focus on cardiac arrest. *Restor Neurol Neurosci* 2003;21:123-139
 46. Rosén H, Rosengren L, Herlitz J, Blomstrand C. Increased serum levels of the S100 protein are associated with hypoxic brain damage after cardiac arrest. *Stroke* 1998;29:473-477
 47. Rosén H, Sunnerhagen KS, Herlitz J, Blomstrand C, Rosengren L. Serum levels of the brain-derived proteins S-100 and NSE predict long-term outcome after cardiac arrest. *Resuscitation* 2001;49:183-191
 48. Böttiger BW, Möbes S, Glätzer R et al. Astroglial protein S-100 is an early and sensitive marker of hypoxic brain damage and outcome after cardiac arrest in humans. *Circulation* 2001;103:2694-2698
 49. Fogel W, Krieger D, Veith M et al. Serum neuron-specific enolase as early predictor of outcome after cardiac arrest. *Crit Care Med* 1997;25:1133-1138
 50. Martens P. Serum neuron-specific enolase as a prognostic marker for irreversible brain damage in comatose cardiac arrest survivors. *Acad Emerg Med* 1996;3:126-131
 51. Wijdicks EF, Hijdra A, Young GB, Bassetti CL, Wiebe S;

- Quality Standards Subcommittee of the American Academy of Neurology. Practice parameter: prediction of outcome in comatose survivors after cardiopulmonary resuscitation (an evidence based review): report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology* 2006;67:203-210
52. Els T, Kassubek J, Kubalek R, Klisch J. Diffusion-weighted MRI during early global cerebral hypoxia: a predictor for clinical outcome? *Acta Neurol Scand* 2004;110:361-367
 53. Torbey MT, Selim M, Knorr J, Bigelow C, Recht L. Quantitative analysis of the loss of distinction between gray and white matter in comatose patients after cardiac arrest. *Stroke* 2000;31:2163-2167
 54. Arbelaez A, Castillo M, Mukherji SK. Diffusion-weighted MR imaging of global cerebral anoxia. *AJNR Am J Neuroradiol* 1999;20:999-1007
 55. Wijdicks EF, Campeau NG, Miller GM. MR imaging in comatose survivors of cardiac resuscitation. *AJNR Am J Neuroradiol* 2001;22:1561-1565
 56. de Crespigny AJ, Röther J, Beaulieu C, Moseley ME, Hoehn M. Rapid monitoring of diffusion, DC potential, and blood oxygenation changes during global ischemia: effects of hypoglycemia, hyperglycemia, and TTX. *Stroke* 1999;30:2212-2222
 57. Fischer M, Bockhorst K, Hoehn-Berlage M, Schmitz B, Hossmann KA. Imaging of the apparent diffusion coefficient for the evaluation of cerebral metabolic recovery after cardiac arrest. *Magn Reson Imaging* 1995;13:781-790
 58. Li F, Silva MD, Liu KF, Helmer KG et al. Secondary decline in apparent diffusion coefficient and neurological outcomes after a short period of focal brain ischemia in rats. *Ann Neurol* 2000;48:236-244
 59. Wijman CA, Mlynash M, Caufield AF et al. Prognostic value of brain diffusion-weighted imaging after cardiac arrest. *Ann Neurol* 2009;65:394-402
 60. Wu O, Sorenson AG, Benner T, Singhal AB, Furie KL, Greer DM. Comatose patients with cardiac arrest: predicting clinical outcome with diffusion-weighted MRI imaging. *Radiology* 2009;252:173-181
 61. Mlynash M, Campbell DM, Leproust EM et al. Temporal and spatial profile of brain diffusion-weighted MRI after cardiac arrest. *Stroke* 2010;41:1665-1672
 62. Laureys S, Goldman S, Phillips C et al. Impaired effective cortical connectivity in vegetative state: preliminary investigation using PET. *Neuroimage* 1999;9:377-382
 63. Marino S, Bramanti P. Neurofunctional imaging in differential diagnosis and evaluation of outcome in vegetative and minimally conscious state. *Funct Neurol* 2009;24:185-188
 64. Luccichenti G, Sabatini U. Colouring rehabilitation with functional neuroimaging. *Funct Neurol* 2009;24:189-193
 65. Di H, Boly M, Weng X, Ledoux D, Laureys S. Neuroimaging activation studies in the vegetative state: predictors of recovery? *Clin Med* 2008;8:502-507
 66. Coleman MR, Davis MH, Rodd JM et al. Towards the routine use of brain imaging to aid the clinical diagnosis of disorders of consciousness. *Brain* 2009;132:2541-2552
 67. Gofton TE, Chouinard PA, Young GB et al. Functional MRI study of the primary somatosensory cortex in comatose survivors of cardiac arrest. *Exp Neurol* 2009;217:320-327
 68. Bassetti C, Bomio F, Mathis J, Hess CW. Early prognosis in coma after cardiac arrest: a prospective clinical, electrophysiological, and biochemical study of 60 patients. *J Neurol Neurosurg Psychiatry* 1996;61:610-615
 69. Sherman AL, Tirschwell DL, Micklesen PJ, Longstreth WT Jr, Robinson LR. Somatosensory potentials, CSF creatine kinase BB activity, and awakening after cardiac arrest. *Neurology* 2000;54:889-894

Finito di stampare
nel mese di maggio 2011
dallo Stab. Tipolit. Ugo Quintily SpA - Roma